

## **CIMPLE PLUS SHAMPOO**

Veko's Cimple shampoo is prescribed to treat pets suffering from skin infections caused by ectoparasites (fleas, tics, lice) and also to control fungal skin infections caused by the ectoparasites or other causes. Generic Equivalent to: **Clinar-M** 

For: Dogs

#### **Benefits:**

- For the treatment of ticks, fleas & mites' infestation of skin.
- Cimple shampoo also heals and prevents any fungal induced infections.
- Along with moisturizing the skin and coat, the shampoo leaves a lovely fragrance, eliminating the bad odour.

#### **How it Works:**

Cypermethrin is a synthetic pyrethroid and a broad-spectrum insecticide, it behaves as a fast-acting neurotoxin in parasites, destructing their nerve tissue. The other ingredient Miconazole Nitrate is imidazole antifungal and works against fungi and yeast such as Malassezia pachydermatis. Miconazole inhibits ergosterol incorporation into cell membranes so increasing concentrations of cytotoxic hydrogen peroxide within the fungal cell wall



200ml Bottle









#### **Precautions:**

- FOR EXTERNAL USE ONLY.
- · Keep out of reach of children.
- · Wash hands after using
- · Avoid contact with eyes and ears
- NOT FOR USE IN CATS

#### What are the side effects:

When used correctly, Cimple shampoo is usually well tolerated and will not cause any side effects. If you do witness any unusual symptoms, report them as soon as possible so that the correct response may be taken. This can include reduced doses or other adjustments to prevent further complications. Emergency medical assistance may also be required in more serious cases. Commonly reported side effects include irritated, flaky, or dry skin.

#### How to use the Product:

Wet the hair coat with warm water and apply sufficient amount of shampoo to create a rich lather. Massage shampoo into wet hair coat, lather freely. Allow to remain on hair for 5 to 10 minutes and then rinse thoroughly with clean water.







# CIMPLE PLUS SHAMPOO

### How Often It should be used?

It can be used initially 3-2 times a week for 4 weeks, then reducing to once a week or as directed by your veterinarian.

Active Ingredient	Amount
Cypermethrin	1 %w/w
Miconazole nitrate	2%w/w

